

Instructions

Errata for ICD-10-AM Tabular List, ICD-10-AM Alphabetic Index, ACHI Tabular List and ACHI Alphabetic Index are produced from a database. This has resulted in the format shown below. Each row represents an item/change to be made and contains the following information necessary to complete the errata item.

Tabular/Index Entry	The Tabular Entry	(Block:code) or Inde	ex Entry (Main term:subterm) t	to which the errata item applies. For example:
	ICD-10-AM Tab Lis	st:R02	ICD-10-AM Tabular List, C	ode R02
	ACHI Tab List:188	6:34533-00	ACHI Tabular List, Block 1	886, Code 34533-00
	ACHI Index:Revision implantable spinal	on:device:infusion,	ACHI Alphabetic Index,	Revision - device
				infusion, implantable spinal
Note Type				s. To use this errata, it is important to think of a ments as shown in the examples below.
Code	R32 Unsp	ecified urinary inco	ontinence —	Code Description
Standard ——	▼ 1808 Enure	esis NOS		Inclusional
	Exclu	udes: nonorganic en	uresis (F98.0)	Excludes
			ence and other specified	Excludes
	Index Entry			
	-			
Term ———	Dacryocystitis	(acute) (phl	egmonous) H04.3	— Code
				NEM
Term ———	- syphilitic A	.52.7† H06.0*		Code
				—— Code
Term	Endoscopy, en	doscopic – s	ee also Panendoscopy	See Also
				NEM
Term	- brain (intra	ventricular)	40903-00 [1]	— Block
				— Code
Term	with biopsy	40903-	01 [12]	Block
				—— Code
Instruction	Type of change to	be made. For exampl	e:	
Tabular/Index Entry		Note Instruction Type	Existing New Value Value	indicates a term to be added at: Complications
ICD-10-AM Index:Complica	tions:umbilical cord	Term Add	velamentous insertion	- umbilical cord velamentous insertion

Existing Value

vascular access

- port, implantable

New Value

Note

Туре

Term

A 'Delete' instruction followed by an asterisk () indicates that there are subterms to be deleted as a

result of this deletion. In the example given above, subterms and notes to be consequently deleted are:

It should be assumed that if a Tabular 'Code' or an Index 'Term' is deleted, all its accompanying notes

('Code Description', 'Include', 'Exclude', 'NEM', 'Code', 'Block', 'See Also', etc) need to be deleted.

Text of the entry before the errata change.

Text with errata in track changes.

Instruction

Delete*

indicates a term to be deleted at:
Revision
- device

- - port, implantable vascular access

- - - for infusion of agents via catheter

- - - - central venous (CVC) 34530-00 [766]

- - - - spinal 39126-00 [56]

---- venous 34530-00 [766]

Eighth Edition - Errata 1, 21 June 2013, for implementation 1 July 2013

ACHI Index:Revision:device:port, implantable

Tabular/Index Entry

vascular access

Existing Value

New Value

ERRATA 1, June 2013

ICD-10-AM/ACHI/ACS Eighth Edition, 2013



e	the rm to which nge applies	Type of change to be made	Text before making the change	Text with errata in track changes
Tabular/Index Entry	Туре	Instruction	Existing Value	New Value
ICD-10-AM Tab List:K81	Excludes	Revise	with cholelithiasis (K80)	with cholelithiasis <u>or</u> <u>choledocholithiasis (</u> K80)
ICD-10-AM Tab List:L55	Includes	Add		burns from exposure to man- made ultraviolet radiation
ICD-10-AM Tab List:L59	Excludes	Revise	radiation burns: see Burn by site	radiation burns: see Burn by site • NOS (T20–T31) • ultraviolet (man-made) (sun) (L55)
ICD-10-AM Tab List:O99.7	Excludes	Revise	pruritis in pregnancy (O26.88)	pruritis-<u>pruritus</u> in pregnancy (O26.88)
ICD-10-AM Tab List:T66	Excludes	Revise	specified adverse effects of radiation, such as: • sunburn (L55)	specified adverse effects of radiation, such as: • sunburn <u>(from man-made</u> <u>ultraviolet radiation)</u> (L55)
ICD-10-AM Tab List:Z06.5	Use additiona l code	Revise	Use additional code (B95–B96) to identify infectious agents resistant to beta-lactam antibiotics.	Use additional code (B95– B96) to identify infectious agents resistant to beta- lactam antibiotics <u>if</u> <u>applicable</u> .
ICD-10-AM Tab List:Z06.6	Use additiona l code	Revise	Use additional code (B95–B96) to identify infectious agents resistant to antibiotics.	Use additional code (B95– B96) to identify infectious agents resistant to antibiotics if applicable.
ICD-10-AM Tab List:Z06.70	Use additiona l code	Revise	Use additional code (B95–B97) to identify infectious agents resistant to antimicrobial drugs.	Use additional code (B95– B97) to identify infectious agents resistant to antimicrobial drugs <u>if</u> <u>applicable</u> .
ICD-10-AM Tab List:Z06.73	Use additiona l code	Revise	Use additional code (B97) to identify infectious agents resistant to antiviral drugs.	Use additional code (B97) to identify infectious agents resistant to antiviral drugs <u>if</u> <u>applicable</u> .
ICD-10-AM Index:Abnormal:abnormality, abnormalities:parturition:mother	See	Revise	— see Delivery/complicated	— see Delivery/complicated Pregnancy/complicated by
ICD-10-AM Index: Anomaly, anomalous: bone: cranium	Code	Revise	Q79.93	Q79.93 <u>Q75.9</u>
ICD-10-AM Index:Anomaly, anomalous:bone:frontal	Code	Revise	Q79.93	Q79.93<u>Q75.9</u>
ICD-10-AM Index:Anomaly, anomalous:bone:head	Code	Revise	Q79.93	Q79.93<u>Q75.9</u>
ICD-10-AM Index:Birth:complications in mother	NEC & Code See	Add Revise	- complications in mother — see Delivery/complicated	- complications in mother <u>NEC</u> — <i>see</i> <i>Delivery/complicated</i> (see <i>also Pregnancy/complicated</i> <i>by</i>) 075.9
ICD-10-AM Index:Burn	NEM	Delete	Burn (radiation)	Burn (radiation)
ICD-10-AM Index:Burn	Term, NEC & Code	Add		- radiation NEC T30.0

	ICD-10-AM Index:Burn:radiation	Term, NEMs & See	Add	- radiation NEC T30.0	- radiation NEC T30.0 ultraviolet (man-made) (rays) (sun) — see Sunburn
	ICD-10-AM Index:Cholangitis:with calculus, bile duct:and cholecystitis	NEMs	Add	and cholecystitis K80.4-	and cholecystitis (acute) (chronic) K80.4-
	ICD-10-AM Index: Choledocholithiasis:with:cholecystitis	NEMs	Add Revise	cholecystitis (and cholangitis) K80.4-	cholecystitis <u>(acute)</u> (<u>chronic)</u> (and <u>with</u> cholangitis) K80.4-
	ICD-10-AM Index:Effect, adverse:ultraviolet:burn	See	Revise	— see Burn	— see Burn<u>Sunburn</u>
	ICD-10-AM Index:Erythema, erythematous:due to:burn:sun	Term & code	Delete	burn sun L55.0	burn
	ICD-10-AM Index:Erythema, erythematous:due to:burn	Term, NEMs & Code	Add	burn	burn ultraviolet radiation (man- made) (rays) (sun) L55.0
	ICD-10-AM Index:Exposure	See also	Revise	(see also Contact/with)	(see also Contact/with)
	ICD-10-AM Index:Fluid:free intraperitoneal:in:pelvic inflammatory disease	See	Revise	— see Disease/diseased/pelvis, pelvic/inflammatory	— see Disease,4 diseased/pelvis, pelvic/inflammatory
	ICD-10-AM Index: Haemorrhoids:strangulated	NEC	Add	- strangulated (see also Haemorrhoids/by degree) K64.8	- strangulated <u>NEC</u> (see also Haemorrhoids/by degree) K64.8
	ICD-10-AM Index: Haemorrhoids:thrombosed	NEC	Add	- thrombosed (see also Haemorrhoids/by degree) K64.8	- thrombosed <u>NEC (see also</u> Haemorrhoids/by degree) K64.8
	ICD-10-AM Index:Haemorrhoids:ulcerated	NEC	Add	- ulcerated (see also Haemorrhoids/by degree) K64.8	- ulcerated <u>NEC (see also</u> Haemorrhoids/by degree) K64.8
	ICD-10-AM Index: Hydrocephalus:congenital:with spina bifida:communicating	Term & Code	Delete	 - congenital - with spina bifida communicating Q03.81 	 congenital - with spina bifida <u>communicating Q03.81</u>
	ICD-10-AM Index: Hydrocephalus:congenital	Term & Code	Add	- congenital	- congenital communicating Q03.81
	ICD-10-AM Index:Hypoplasia:leg	NEM & Code NEC	Revise Add	- leg (lower) Q72.81	- leg (lower congential) <u>NEC</u> Q72.81 Q72.89
	ICD-10-AM Index:Hypoplasia:leg:meaning lower limb	Term See Code	Revise Delete Add	meaning lower limb — see Hypoplasia/limb/lower	meaning- lower limb see Hypoplasia/limb/lower Q72.81
	ICD-10-AM Index:Injury:obstetric	See also	Revise	(see also Delivery/complicated by/injury by type)	(see also Delivery/complicated by/injury by type Injury/instrumental/obstetric)
	ICD-10-AM Index:Injury:ultraviolet rays	Term NEMs	Revise Add	- ultraviolet rays NEC T66	- ultraviolet rays -(radiation) (rays) NEC T66
	ICD-10-AM Index:Injury:ultraviolet	Term & See	Add Add		<u> burn — see Sunburn</u>
	ICD-10-AM Index:Lymphoma:Hodgkin	Term, NEC & Codes	Add	- Hodgkin	- Hodgkin classical NEC (M9650/3) <u>C81.7</u>

	ICD-10-AM Index:Melanosis:precancerous	See also	Revise	— see also Melanoma/in-situ	— see also Melanoma/insitu
	ICD-10-AM Index:Neoplasm, neoplastic:alveolar:ridge or process	The sign◆	Delete	Neoplasm, neoplastic - alveolar - ridge or process ◆	Neoplasm, neoplastic - alveolar ridge or process ◆
	ICD-10-AM Index:Neoplasm, neoplastic: alveolar:ridge or process:lower	The sign◆	Delete	Neoplasm, neoplastic - alveolar - ridge or process lower ◆	Neoplasm, neoplastic - alveolar ridge or process lower
	ICD-10-AM Index:Neoplasm, neoplastic: alveolar:ridge or process:upper	The sign∳	Delete	Neoplasm, neoplastic - alveolar - ridge or process upper ◆	Neoplasm, neoplastic - alveolar ridge or process upper
	ICD-10-AM Index:Neoplasm, neoplastic:jaw:bone	The sign♠	Delete	Neoplasm, neoplastic - jaw bone ◆ lower ◆ upper ◆	Neoplasm, neoplastic - jaw bone lower upper
	ICD-10-AM Index:Neoplasm, neoplastic:malar	The sign◆	Delete	Neoplasm, neoplastic - malar ◆	Neoplasm, neoplastic - malar 🔶
	ICD-10-AM Index:Neoplasm, neoplastic:mandible	The sign◆	Delete	Neoplasm, neoplastic - mandible ◆	Neoplasm, neoplastic - mandible ✦
1	ICD-10-AM Index:Neoplasm, neoplastic:mandible:alveolar:ridge or process	The sign♠	Delete	Neoplasm, neoplastic - mandible - alveolar ridge or process ◆	Neoplasm, neoplastic - mandible alveolar ridge or process 4
	ICD-10-AM Index:Neoplasm, neoplastic:marrow	The sign◆	Delete	Neoplasm, neoplastic - marrow ◆	Neoplasm, neoplastic - marrow
	ICD-10-AM Index:Neoplasm, neoplastic:maxilla, maxillary	The sign◆	Delete	Neoplasm, neoplastic - maxilla, maxillary ◆	Neoplasm, neoplastic - maxilla, maxillary ◆
1	ICD-10-AM Index:Neoplasm, neoplastic:maxilla, maxillary:alveolar:ridge or process	The sign◆	Delete	Neoplasm, neoplastic - maxilla, maxillary alveolar ridge or process ◆	Neoplasm, neoplastic - maxilla, maxillary alveolar ridge or process
	ICD-10-AM Index:Neoplasm, neoplastic:vomer	The sign◆	Delete	Neoplasm, neoplastic - vomer ◆	Neoplasm, neoplastic - vomer
1	ICD-10-AM Index:Nephritis, nephritic:complicating pregnancy, childbirth or puerperium	Term	Revise	- complicating pregnancy, childbirth or puerperium O26.81	- complicating pregnancy , childbirth or puerperium O26.81
	ICD-10-AM Index: Tumour:malignant:spindle cell	See also	Add	spindle cell (type) (M8004/3) C80	spindle cell (type) (M8004/3) <u>(see also</u> <u>Carcinoma/spindle cell)</u> C80
	ICD-10-AM Index:Tumour:spindle:cell	Morph code	Revise	cell (M8032/1) benign (M8032/0) malignant (M8032/3) —see Carcinoma/spindle cell	cell (<u>M8032/1M8004/1</u>) benign (<u>M8032/0M8004/0</u>) malignant (<u>M8032/3M8004/3</u>) —see <u>also</u> Carcinoma/spindle cell

ICD-10-AM Index:EXT:Cut, cutting	NEC & Code	Add and Revise	Cut, cutting (accidental) (any part of body) (by) (see also <i>Contact/with/by object or</i> <i>machine</i>) W49	Cut, cutting (accidental) (any part of body) (by) <u>NEC</u> (see also Contact/with/by <u>type of</u> object or machine) <u>W49W45.9</u>
ICD-10-AM Index:EXT:Piercing	NEC & Code	Add and Revise	Piercing (see also Contact/with/by type of object or machine) W49	Piercing <u>NEC</u> (see also Contact/with/by type of object or machine) <u>W49</u> <u>W45.9</u>
ICD-10-AM Index:EXT:Puncture, puncturing	NEC & Code	Add and Revise	Puncture, puncturing (see also Contact/with/by type of object or machine) W49	Puncture, puncturing <u>NEC</u> (see also Contact/with/by type of object or machine) W49 <u>W45.9</u>
ACHI Tab List:865	Excludes	Revise	that by laparoscopic approach (30532-04 [854], 30532-05, 30533 [863])	that by laparoscopic approach (30532-04-<u>30532-00</u> [854], 30532-05, 30533 [863])
ACHI Tab List:911:32093-00	Includes	Revise	polypectomy by or using: • biopolar cautery	polypectomy by or using: • biopolar <u>bipolar</u> c autery
ACHI Index:Arthroplasty:knee:for joint replacement:total	NEM	Add	total (unilateral) 49518-00 [1518]	total (unilateral) <u>(with</u> <u>patellofemoral joint</u> <u>replacement)</u> 49518-00 [1518]
ACHI Index:Arthroplasty:knee:for joint replacement:total:patellofemoral joint	Term, Code & Block	Delete	patellofemoral joint 49534-01 [1518]	
ACHI Index:Banding:gastric, for obesity	NEC, Code & Block	Add	- gastric, for obesity (with replacement)	- gastric, for obesity (with replacement) <u>NEC 30511-</u> 04 [889]
ACHI Index:Destruction:scar, by laser	See	Revise	— see Resurfacing/scar/by site	— see Resurfacing/ scar <u>skin</u> /by site
ACHI Index:Gastrectomy:sleeve:for procedures other than obesity	Term	Revise	for procedures other than obesity 30523-00 [879]	for procedures other than <u>for</u> obesity 30523-00 [879]
ACHI Index:Replacement:joint:knee:total	NEM	Add	total	total <u>(with patellofemoral</u> joint replacement)
ACHI Index: Replacement:joint:knee:total:bilateral:patell ofemoral joint	Term, Code & Block	Delete	patellofemoral joint 49534-01 [1518]	patellofemoral joint 49534 01 [1518]
ACHI Index: Replacement:joint:knee:total:unilateral:patel lofemoral joint	Term,	Delete	patellofemoral joint 49534-01 [1518]	patellofemoral joint 49534-01 [1518]
ACHI Index:Resurfacing:patellofemoral	Terms	Add	- patellofemoral 49534-01 [1518]	- patellofemoral 49534-01 [1518] with total knee replacement (unilateral) 49518-00 [1518] bilateral 49519-00 [1518]

AUSTRALIAN CODING STANDARDS

Pg 15 GUIDE FOR USE Revise 5. For combination codes (see ACS 0015 Combination codes) where a diagnosis When and AM code describes multiple concepts (ie a combination code) and any concept within the criteria of COF 1, and is not represented by another code with a COF 1 value, then assign combination code (see Example 2). Pg 18 EXAMPLE 11: Patient admitted with uncontrolled Type 2 diabetes. The patient is known to have nephrop 3) but no other complications of the diabetes. During admission the patient develops acute 2 – Type 2 diabetes mellitus uncontrolled 1 – Acute kidney failure 24 – Type 2 diabetes mellitus with acute kidney failure 2 – Type 2 diabetes mellitus with chronic kidney disease	eat code meets the a COF 1 <u>to the</u> pathy (CKD stage
Revise AM code describes multiple concepts (ie a combination code) and any concept within the criteria of COF 1, and is not represented by another code with a COF 1 value, then assign combination code (see Example 2). Pg 18 EXAMPLE 11: Patient admitted with uncontrolled Type 2 diabetes. The patient is known to have nephrop 3) but no other complications of the diabetes. During admission the patient develops acute 2 – Type 2 diabetes mellitus uncontrolled 1 – Acute kidney failure Pariate 24 – Type 2 diabetes mellitus with acute kidney failure	eat code meets the a COF 1 <u>to the</u> pathy (CKD stage
Pg 18 Patient admitted with uncontrolled Type 2 diabetes. The patient is known to have nephrop 3) but no other complications of the diabetes. During admission the patient develops acute 2 – Type 2 diabetes mellitus uncontrolled 1 – Acute kidney failure 24 – Type 2 diabetes mellitus with acute kidney failure	
1 – Acute kidney failure 21 – Type 2 diabetes mellitus with acute kidney failure	
2 - Chronic kidney disease, stage 3	
0102 HIV/AIDS	
<i>Pg 42</i> EXAMPLE 1:	
Revise Antiretroviral drug induced haemolytic anaemia with HIV positive status. A patient is admitted with oral candidiasis due to HIV infection.	
Assign and sequence as:	
Oral candidiasis HIV disease B20 <i>Candidal stomatitis</i> HIV disease B20 <i>Human immunodeficiency virus [HIV] disease result</i> <i>infectious and parasitic diseases</i>	ting in
Pg 43 EXAMPLE 2: A patient with AIDS is admitted with a principal diagnosis of Kaposi sarcoma of skin, an diagnoses of wasting syndrome and CMV retinitis.	nd additional
Assign and sequence as:	
Revise Kaposi sarcoma C46.0 <u>Kaposi sarcoma of skin</u> Drug induced nonautoimmus	me haemolytic
HIV disease M9140/3 Kaposi sarcoma B21 Human immunodeficiency virus [HIV] disease result malignant neoplasmsAntiviral drugs causing adversed	
therapeutic useCMV retinitisH30.9Chorioretinal inflammation, unspecifiedB25.8Other cytomegaloviral diseases	
B25.8Other cytomegaloviral diseasesWasting syndromeR64Cachexia	
0402 CYSTIC FIBROSIS	
Pg 79 Cystic fibrosis (CF) should be coded with the appropriate code from E84 Cystic fibrosis code(s) for any specified manifestation(s), irrespective of whether the manifestation(s) manifestation and diagnoses. More than one code from E84 Cystic fibrosis should be used if the presents with multiple manifestations of CF.	neets ACS 0002

1615 SPECIFIC INTERVENTIONS FOR THE SICK NEONATE

Pg 186

Revise

2. CODE THE FOLLOWING INTERVENTION WHEN PERFORMED:

Parenteral fluid therapy

96199-07 [1920] Intravenous administration of pharmacological agent, nutritional substance and 96199-08 [1920] Intravenous administration of pharmacological agent, electrolytes should be assigned when used for management of carbohydrate, hydration or electrolyte disorders.

Examples of diagnoses that may require parenteral fluid therapy are 'neonatal hypoglycaemia' (P70.-*Transitory disorders of carbohydrate metabolism specific to fetus and newborn*) or 'other transient neonatal electrolyte disorders' (P74.- *Other transitory neonatal electrolyte and metabolic disturbances*).

Note: These codes should **not be assigned** when these procedures are **part of resuscitation at birth**.

ICD-10-AM TABULAR LIST OF DISEASES

Page 1

CHAPTER 1 CERTAIN INFECTIOUS AND PARASITIC DISEASES (A00-B99)

Includes: diseases generally recognised as communicable or transmissible

Use additional code (Z06-Z07) to identify resistance to antimicrobial and antineoplastic drugs

Excludes: carrier or suspected carrier of infectious disease (Z22.-) certain localised infections - see body system-related chapters infectious and parasitic diseases complicating pregnancy, childbirth and the puerperium [except obstetrical tetanus] (098.-)

> infectious and parasitic diseases specific to the perinatal period [except tetanus neonatorum, congenital syphilis, perinatal gonococcal infection and perinatal human immunodeficiency virus [HIV] disease] (P35–P39) influenza and other acute respiratory infections (J00–J22)

PAGE 32

CHAPTER 2 NEOPLASMS (C00–D48)

Note:

...

8. Use of the third edition of International Classification of Diseases for Oncology (ICD-O)


For certain morphological types, Chapter 2 provides a rather restricted topographical classification, or none at all. The topography codes of ICD-O use for all neoplasms essentially the same three- and four- character categories that Chapter 2 uses for malignant neoplasms (C00-C77, C80.-), thus providing increased specificity of site for other neoplasms (malignant secondary (metastatic), benign, in situ and uncertain or unknown).

It is therefore recommended that agencies interested in identifying both the site and morphology of tumours, e.g. cancer registries, cancer hospitals, pathology departments and other agencies specialising in cancer, use ICD-O.

Use additional code (Z07) to identify resistance to antineoplastic drugs



	<i>Excludes:</i> abnormal white blood cells (count) (R72) basophilia (D75.8) immunity disorders (D80–D89) neutropenia (D70) preleukaemia (syndrome) (D46.9)
072.0	Genetic anomalies of leukocytes Anomaly (granulation)(granulocyte) or syndrome: • Alder • May-Hegglin • Pelger-Huët Hereditary: Hereditary: • leukocytic: • hypersegmentation • hyposegmentation • leukomelanopathy Excludes: Chediak(-Steinbrinck)-Higashi syndrome (E70.3)





CHAPTER 19 INJURY, POISONING AND CERTAIN OTHER **CONSEQUENCES OF EXTERNAL CAUSES** (S00–T98)

. . .

. 1 1

Fracture including:
fracture:
closed:
 comminuted
 depressed
• elevated
• fissured
 greenstick
• impacted
• linear
• march
• simple
 slipped epiphysis
• spiral
• dislocated
. diamla and

displaced

PAGE 504

©Z07

Resistance to antineoplastic drugs

}

Note: Assign Z07 as an additional code to identify resistance to antineoplastic drugs in the treatment of conditions classified elsewhere in Chapter 2.

with or without delayed healing



ICD-10-AM

ALPHABETIC INDEX OF DISEASES

PAGE 21

Admission (for)

- breast

- - augmentation or reduction Z41.1
- - removal, prophylactic for risk-factors related to malignant neoplasms Z40.00
- ... - prophylactic organ removal surgery Z40.9
- - organ removal for risk-factors related to malignant neoplasms - - - breast Z40.00
- - ovary Z40.01
- - specified organ NEC Z40.08
- - specified NEC Z40.8
- psychotherapy NEC Z50.4

PAGE 115

Diabetes, diabetic (controlled) (mellitus) (without complication) E1-.9

Note:

- The three character subdivision for diabetes mellitus is: 0 Type 1 (IDDM) 1 Type 2 (NIDDM) 3 Other specified 4 Unspecified
- for stabilisation E1-.65
- with
- - abnormal sweating (gustatory) E1-.43
- ... - - ulcer, skin E1-.69
- - foot (see also ACS 0401 Diabetes mellitus and intermediate hyperglycaemia/6 Diabetic foot) E1-.69
- - lower extremity E1-.69
- -- unspecified complication E1-.8
- - vitreous haemorrhage E1-.33



Hyperglycaemia, hyperglycaemic R73

- with diabetes (mellitus) see Diabetes, diabetic
- coma E1-.02
- intermediate (without complication) E09.9
- - with
- - acanthosis nigricans E09.72
- ...
- - tubulo-interstitial changes E09.21
- --- unspecified complication E09.8
- - in pregnancy, childbirth or puerperium O24.5-

PAGE 318

	Malig	gnant			Uncertain or unknown
Neoplasm, neoplastic	Primary C80.9	Secondary C79.9	In situ D09.9	Benign D36.9	behaviour D48.9
- Santorini's duct	C25.3	C78.8	D01.7	D13.6	D37.7 <u>1</u>

PAGE 372

Prophylactic

- organ removal (for neoplasm management) for risk-factors related to malignant neoplasms
- - breast Z40.00
- - ovary Z40.01
- - specified NEC Z40.08
- pharmacotherapy Z29.2
- •••

- surgery (see also Surgery/prophylactic) Z40.9

- - specified NEC Z40.8

PAGE 384

Removal (from) (of) - continued

•••

- organ, prophylactic (for neoplasm management) for risk-factors related to malignant neoplasms
- - breast Z40.00
- - ovary Z40.01
- - specified organ NEC Z40.08
- pin Z47.0





Surgery


- prophylactic Z40.9
- - organ removal for risk-factors related to malignant neoplasms (for neoplasm management)
- - breast Z40.00
- - ovary Z40.01
- - specified organ NEC Z40.08
- - specified NEC Z40.8
- reconstructive (following healed injury or operation) Z42.9

ACHI TABULAR LIST OF INTERVENTIONS

PAGE 82-83

623	Replacement of aortic valve
	 <i>Code also when performed:</i> • cardiopulmonary bypass (38600-00, 38603-00 [642]) • operative management of acute infective endocarditis during heart valve procedure (38493-00 [666])
38488-08	Percutaneous replacement of aortic valve with bioprosthesis Transcatheter aortic valve implantation
	Includes: cardiac catheterisation valvuloplasty
	Code also when performed: • coronary angiography (38215-00, 38218 [668])

PAGE 83





634	Replacement of tricuspid valve
	 Code also when performed: • cardiopulmonary bypass (38600-00, 38603-00 [642]) • operative management of acute infective endocarditis during heart valve procedure (38493-00 [666])
38488-10	Percutaneous replacement of tricuspid valve with bioprosthesis
	Includes: cardiac catheterisation valvuloplasty
	Code also when performed: • coronary angiography (38215-00, 38218 [668])

PAGE 84-<u>85</u>

637 Repair or replacement of pulmonary valve

38270-03 Percutaneous balloon pulmonary valvuloplasty

Includes: cardiac catheterisation

38488-11 Percutaneous replacement of pulmonary valve with bioprosthesis Transcatheter pulmonary valve implantation

> Includes: cardiac catheterisation valvuloplasty

Code also when performed: • coronary angiography (38215-00, 38218 **[668]**)

PAGE 110

 738
 Venous catheterisation

 ∇ 0042
 ...

 13300-00
 Catheterisation/cannulation of other vein in neonate

 ∇ 1615
 Includes: infusion



. . .

1889 Other therapeutic interventions on respiratory system

92210-00 Nitric oxide therapy ∇ 1615

> Code first: • ventilatory support for neonates (13882 [569], 92209 [570])

ACHI ALPHABETIC INDEX OF INTERVENTIONS

PAGE 49

Correction — see also Repair

- ... - ptosis
- - breast 45556-00 [1755]
- - eyebrow (bilateral) 45588-01 [1675]
- - unilateral 45587-01 [1675]



PO Box 170

t: +61 2 9351 9772 f: +61 2 9351 9603 Lidcombe NSW 1825 e: enquiries@accd.net.au w: accd.net.au

AUSTRALIAN CODING STANDARDS FOR ICD-10-AM AND ACHI

PAGE 186

1615 SPECIFIC INTERVENTIONS FOR THE SICK NEONATE

•••

2. CODE THE FOLLOWING INTERVENTION WHEN PERFORMED:

•••

Catheterisation in a neonate

1 <u>3300-00 [738]</u>	Catheterisation/cannulation of other vein in neonate
13300-01 [738]	Scalp vein catheterisation/cannulation in neonate
13300-02 [738]	Umbilical vein catheterisation/cannulation in neonate
13319-00 [738]	Central vein catheterisation in neonate
13303-00 [694]	Umbilical artery catheterisation/cannulation in neonate
Note: When m	ultiple catheterisations are performed during an episode of care and the same
procedure code a	polies, assign the procedure code once only.

STANDARDS INDEX

PAGE 250			
C			
 Cystic fibrosis 0402 , 34 - <u>79</u>			



Eighth Edition - Errata 2, 15 December 2013, for implementation 1 January 2014