



IHACPA

Independent Health and Aged Care Pricing Authority

**ICD-10-AM/ACHI/ACS
Twelfth Edition**

Errata 4

For implementation

1 April 2023

How to use this document

The ICD-10-AM/ACHI/ACS classification system is used for classifying admitted patient care and comprises the following:

- International Statistical Classification of Diseases and Related Health Problems, Tenth Revision, Australian Modification (ICD-10-AM)
- Australian Classification of Health Interventions (ACHI)
- Australian Coding Standards (ACS).

For the published hardcopy books, ICD-10-AM/ACHI/ACS errata may be issued to correct errors and typographical issues that have been identified since the time of publishing. Users are required to incorporate these corrections when using the published hardcopy books.

This document lists corrections to be made to each of the hardcopy books listed in the following order:

- ICD-10-AM Tabular List
- ICD-10-AM Alphabetic Index
- ACHI Tabular List
- ACHI Alphabetic Index
- Australian Coding Standards

Each correction includes:

- The hardcopy book with a page reference
- Instructions to correct codes or terms (eg add, delete, amend)
- The corrected text without track changes detailed in a grey box
- The existing text with track changes detailed in a white box (eg [added text](#), ~~deleted text~~)

Example of correction

Page #	Add/Delete/Amend <i>Instructional note/code(s)/term(s)</i> , and Add/Delete/Amend <i>Instructional note/code(s)/term(s)</i> <i>Corrected text.</i>
CODE ▽ 0000	Code description <i>Incorrect</i> <i>Corrected</i> text .

ICD-10-AM Tabular List

Page 405	Amend typographical error <i>Note:</i> Munchhausen's (syndrome) by proxy is a disorder in which a carer (usually a parent) fabricates symptoms or induces an illness in the patient (usually a child).
T74.1	Physical abuse Battered: <ul style="list-style-type: none">• baby or child syndrome NOS• spouse syndrome NOS Munchhausen's (syndrome) by proxy <i>Note:</i> Munchhausen's (syndrome) by proxy is a disorder in which a carer (usually a parent) fabricates symptoms or induces and <u>an</u> illness in the patient (usually a child). <i>Use additional external cause code (Y07.0-) to identify perpetrator</i>

ICD-10-AM Alphabetic Index

Page 336	Delete nonessential modifier (NEM), and Add NEM - amoxicillin-clavulanic acid (Augmentin) Z14.01
	Resistance, resistant (to) - amoxicillin, amoxycillin (Augmentin) Z14.11 - amoxicillin-clavulanic acid (Augmentin) Z14.01
Page 337	Amend morphology code - acute infantile (M9751/3) C96.0
	Reticuloendotheliosis - acute infantile (M9754/3 M9751/3) C96.0
Page 337	Amend morphology code - acute, of infancy (M9751/3) C96.0
	Reticulosis (skin) - acute, of infancy (M9754/3 M9751/3) C96.0

ACHI Tabular List

(This section intentionally left blank)

ACHI Alphabetic Index

(This section intentionally left blank)

Australian Coding Standards (ACS)

Page 83	Delete code, block and code title, and Add code, block and code title
<i>37216-00 [1163] Needle biopsy of prostate or seminal vesicle via transrectal route</i>	
0233	MORPHOLOGY
...	
EXAMPLE 2:	
Patient was admitted for a TRUS (transrectal ultrasound guided) biopsy of the prostate under local anaesthesia. Histopathology reported adenocarcinoma (M8140/3) and high grade glandular intraepithelial neoplasia (M8148/2) of the prostate.	
Codes:	C61 <i>Malignant neoplasm of prostate</i>
	M8140/3 <i>Adenocarcinoma NOS</i>
	37218-00 [1163] <i>Needle biopsy of prostate or seminal vesicle</i>
	<u>37216-00 [1163] <i>Needle biopsy of prostate or seminal vesicle via transrectal route</i></u>
Note: M8140/3 <i>Adenocarcinoma NOS</i> is more invasive than M8148/2 <i>Glandular intraepithelial neoplasia, high grade</i> and therefore the morphology for adenocarcinoma /3 is assigned.	

Resequence ICD-10-AM codes, and**Add cross reference to the Note**

Codes:	Z08.9	<i>Follow-up examination after unspecified treatment for malignant neoplasm</i>
	D09.0	<i>Carcinoma in situ, bladder</i>
	M8130/2	<i>Papillary urothelial carcinoma, noninvasive</i>
	...	

Note: For classification purposes, this is **not** considered a recurrence as the morphology is different to the original neoplasm (see ACS 0052 *Same-day endoscopy – surveillance*).

0237**RECURRENCE OF MALIGNANCY**

...

EXAMPLE 4:

Patient admitted for a check cystoscopy under GA, for a previous transitional cell carcinoma (TCC) (M8120/3) of the bladder. A noninvasive papillary TCC (M8130/2) of the bladder was confirmed by biopsy.

Codes:	<u>Z08.9</u>	<u><i>Follow-up examination after unspecified treatment for malignant neoplasm</i></u>
	D09.0	<i>Carcinoma in situ, bladder</i>
	M8130/2	<i>Papillary urothelial carcinoma, noninvasive</i>
	Z08.9	<i>Follow-up examination after unspecified treatment for malignant neoplasm</i>
	36836-00 [1098]	<i>Endoscopic biopsy of bladder</i>
	92514-99 [1910]	<i>General anaesthesia, ASA 99</i>

Note: For classification purposes, this is **not** considered a recurrence as the morphology is different to the original neoplasm (see ACS 0052 *Same-day endoscopy – surveillance*).

Amend block

90469-01 [1338] *Failed vacuum assisted delivery*

1505**DELIVERY AND ASSISTED DELIVERY CODES**

...

EXAMPLE 2:

Patient admitted in labour (39/40). Vacuum delivery attempted but failed, followed by vaginal delivery of healthy infant.

Codes:	O83	<i>Other assisted single delivery</i>
	O66.5	<i>Failed application of vacuum extractor and forceps, unspecified</i>
	Z37.0	<i>Single live birth</i>
	90477-02 [1339]	<i>Assisted vertex delivery</i>
	90469-01 [1343 1338]	<i>Failed vacuum assisted delivery</i>